



Clamp, P. J., De-Loyde, K., Maw, A. R., Gregory, S., Golding, J., & Hall, A. (2020). Factors associated with the development of paediatric chronic otitis media by age 9: A prospective longitudinal cohort study of 6560 children. *Journal of Laryngology and Otology*, 134(11), 998 - 1009. <https://doi.org/10.1017/S0022215120002182>

Peer reviewed version

Link to published version (if available):  
[10.1017/S0022215120002182](https://doi.org/10.1017/S0022215120002182)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via University of Cambridge Press at <https://doi.org/10.1017/S0022215120002182>. Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

**Factors associated with the development of paediatric chronic otitis media by age 9 - A prospective longitudinal cohort study of 6560 children.**

Mr Philip J Clamp <sup>1,2</sup>

Miss Katie De-Loyde <sup>2</sup>

Mr A Richard Maw <sup>2</sup>

Mr Steven Gregory <sup>2</sup>

Prof Jean Golding <sup>2</sup>

Dr Amanda Hall <sup>3,4</sup>

1. Department of Ear, Nose and Throat Surgery, University Hospitals Bristol NHSFT, Bristol, UK.
2. School of Social and Community Medicine, University of Bristol, Bristol UK.
3. Children's Hearing Centre, University Hospitals Bristol NHSFT, Bristol UK
4. Life and Health Sciences, Aston University, Birmingham, UK

**Presentation**

This paper was presented at BACO International, Manchester UK, July 2018.

**Author responsible for corrections and proofs**

Mr Philip J Clamp

Consultant in Ear, Nose and Throat Surgery

St Michael Hospital, University Hospitals Bristol

Southwell Street

Bristol, UK

BS2 8EG

philip.clamp@uhbristol.nhs.uk

0117 9230000

### **Twitter**

@PhilipClampENT

### **Conflicts of interest**

None to declare

### **Funding**

The UK Medical Research Council and the Wellcome Trust (**Grant ref: 102215/2/13/2**) and the University of Bristol currently provide core support for ALSPAC.

Funding for the analyses in this paper were generously provided though a research grant from the “Above and Beyond” charity for central Bristol Hospitals.

## **Abstract**

**Objective:** To study social, health and environmental factors associated with the development of chronic otitis media (COM) by age 9.

**Method:** Prospective, longitudinal, birth cohort study of 6560 children, reviewed at age 9. COM defined as previous surgical history or video-otoscopic changes of tympanic membrane retraction, perforation or cholesteatoma. Non-affected children used control group.

**Results:** Univariate analysis demonstrated an association between COM and otorrhea, snoring, grommet insertion, adenoidectomy, tonsillectomy, hearing loss, abnormal tympanograms and preterm birth. Multivariate analysis suggests many of these factors may be interrelated.

**Conclusion:** The association between COM and otorrhea, abnormal tympanograms and grommets supports the role of the Eustachian tube, otitis media (with effusion or acute) in the pathogenesis of COM. The role of snoring, adenoidectomy and tonsillectomy is unclear. Associations suggested by previous studies (sex, socioeconomic group, parental smoking, maternal education, childcare, crowding, siblings) were not found to be significant predictors in this analysis.

**Keywords:** Otitis media; Cholesteatoma; Tympanic membrane perforation; Paediatrics; Risk factors; Cohort studies; Hearing loss; Otologic Surgical Procedures; Middle ear ventilation; Snoring; Tonsillectomy; Adenoidectomy

## **Introduction**

Otitis media (OM) is a broad term used to describe an inflammatory process within the middle ear cleft. OM, in all its forms, represents a major worldwide health burden<sup>1</sup>. Chronic otitis media (COM) is characterised by more long-term changes to the shape and function of the ear-drum, including tympanic membrane (TM) retraction, perforation and cholesteatoma. COM is a leading cause of permanent hearing loss, especially within developing countries<sup>2-4</sup>. It is associated with reduced quality of life and may require complex ear surgery to remove diseased tissue and preserve or reconstruct hearing function. The diagnosis of COM increases the risk of serious and potentially life-threatening complication including mastoiditis, meningitis, intracranial venous sinus thrombosis and intracranial abscess formation<sup>5</sup>.

Whilst acute otitis media (AOM) and otitis media with effusion (OME) are common in childhood, COM is relatively rare<sup>1</sup>. In the United Kingdom (UK), the prevalence, in British adults, of active and inactive middle ear disease is around 1.5% and 2.6% respectively<sup>6</sup>. It is unclear why some children with a propensity for AOM or OME progress to COM, whilst others do not. Numerous studies have attempted to identify risk factors for COM<sup>6-18</sup>. These are usually case-control studies based on single-point observational studies of affected groups. To date, there have been limited prospectively collected, population studies of the risk factors for COM in children<sup>9, 19</sup>.

The Avon Longitudinal Study of Parents and Children (ALSPAC) is a unique, prospective UK birth cohort study following the health and development of over 14,000 children from before birth. As part of the study, some children underwent video-otoscopic examination their tympanic membranes at age 9 years. A previous study has reviewed these images and reported the prevalence of COM changes including TM retraction, perforation and undiagnosed cholesteatoma<sup>20</sup>. Retraction of the TM pars flaccida and pars tensa was seen in 9.6% and 7.9% of cases respectively. There were 15 cases of suspected cholesteatoma.

The aim of this study is to identify associations and potential risk factors for the development of paediatric COM at age 9. The prospective cohort design allows factors associated with chronic otitis media to be identified without recall bias.

## **Materials and Method**

### *The Avon Longitudinal Study of Parents and Children (ALSPAC).*

ALSPAC or “Children of the 90s study” recruited 14,541 pregnant mothers resident in a defined area (Avon in the southwest of the UK) and with expected date of delivery between 1<sup>st</sup> April 1991 and 31<sup>st</sup> December 1992. This prospective, ethically approved, longitudinal birth cohort study collected health, social, environmental and genetic data for the families and children enrolled. Mothers and their children were followed up using a variety of methods including self-completion questionnaires and direct examination<sup>21</sup>. The ALSPAC website contains details of all the data that is available through a fully searchable data dictionary (<http://www.bris.ac.uk/alspac/researchers/data-access/data-dictionary/>).

Ethical approval for this study was granted by the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees.

### *Study sample*

The study sample consisted of children within ALSPAC who attended a 9-year-old review “Focus clinic” with video-otoscopic data available. Children with a history of cleft lip and/or palate (CLP) were excluded. CLP is known to be a strong independent risk factor for the development of COM and further analysis of the associations/risk factors for these children was deemed unlikely to be informative<sup>22, 23</sup>.

### *Outcome data*

#### *Chronic Otitis Media*

At the 9-year-old Focus clinic, data collected included video-otoscopic imaging of the child's TM and a parental interview regarding the child's Ear, Nose and Throat (ENT) surgical history. This study utilises the method of obtaining, coding and staging the TM images described in a previous paper<sup>20</sup>. For the purposes of this study, COM was defined and categorised as:

- 1) Significant COM: Video-otoscopic evidence of stage two retraction of the pars tensa or flaccida, perforation, cholesteatoma/attic crust, or history of surgery for cholesteatoma or perforation.
- 2) Mild COM: Video-otoscopic evidence of stage one retraction of the pars tensa or flaccida.
- 3) No COM: Video-otoscopic evidence of normal tympanic membrane, with or without middle ear effusion, ventilation tube or tympanosclerosis.

Where an image of the TM was available for one ear only, the outcome of that child was categorised according to that ear only. This was deemed appropriate due to the fact that, of those children with data for both ears available, if no COM was recorded in one ear, 94% of those children also had no COM recorded in the other ear. This system of classification is consistent with previous classification systems used for COM<sup>24</sup>.

#### *ENT and hearing factors*

Information on the child's ENT symptoms was collected by maternal questionnaire at ages 6, 18, 30, 42, 57, 69, 81, 91 and 103 months. Symptoms reported at each time point included: pus/mucus (not wax) leaking from ears, cough for at least two days, having a fever with a cough, snoring for more than a minute at a time and breath holding while asleep.

The following variables were derived from the longitudinal questionnaire data: ear pus/mucus leakage; breath holding and snoring; early snoring, prolonged snoring; cough count, fever with a cough. All of these variables are defined in more detail in the relevant table legend.

At the 9-year-old Focus clinic, information was gathered about the child's history of ENT surgery including: ear surgery for COM; insertion of grommets (categorised as none, once or more than once); history of adenoidectomy; history of tonsillectomy.

Information about the child's hearing and middle ear function were available from the 7-year-old Focus clinic.

### *Child, maternal and socioeconomic factors*

Mother's smoking in pregnancy was obtained from maternal questionnaires at 18 and 32 weeks gestational age. A variable of any smoking in pregnancy was derived. The birthweight and gestational age of the child (categorised as <37 weeks or  $\geq 37$  weeks) were obtained from the maternity records.

Mother's highest education level, the mother's age when the child was born and the parents' occupation, were all collected during pregnancy. Definitions for these variables can be seen in Table 1.

Child factors included the sex of the child and ethnicity, categorised as white or non-white.

Mothers reported how they fed their baby when the child was age 4 weeks, 6 and 15 months of age and variables were derived for: any breastfeeding up to 15 months; duration of breastfeeding. Information on dummy (also known as pacifier, soother or comforter) use was reported in maternal questionnaires when the child was age 4 weeks, 6 and 24 months. A variable of any dummy/pacifier use was derived. Mother's history of grommets was obtained by parental questionnaire.

### *Additional medical factors*

Information on allergy was obtained using skin prick test responses carried out at the age 7 Focus clinic<sup>25</sup>. Atopic dermatitis was measured according to the ISAAC 1994 protocol at the 7-year-old Focus clinic<sup>26-28</sup>. A diagnosis of asthma by a doctor was obtained from maternal questionnaires when the child was age 91 months. All variables relating to medical factors are defined in the relevant Table legends.



### *Environmental factors*

Information about day care was available from parent questionnaires at age 15, 24 and 38 months. A variable was derived of regular day care at any time up to 38 months; any day care up to 38 months and day care of more than 10 hours per week at 38 months.

Information about household crowding was obtained when the child was age 8, 33 and 81 months. Parents were asked about the number of people living in the household (including the number of children) and the number of rooms. The crowding index was calculated (definition in table legend). Parents provided information about damp and mould in the home when the child was aged eight months. At 6, 38 and 54 months, parents reported about the child's exposure to smoking and a variable was derived of any exposure to smoke.

### *Statistical analysis*

Missing data were imputed for the study sample. Multiple imputation was used to impute missing data for all variables of interest. Five imputations were used with 50 iterations.

The Little's MCAR (missing completely at random) test indicated that the data was not missing at random ( $p < 0.001$ ). Data was imputed, but for robustness we also completed all analysis on only the original dataset. This analysis confirms that imputing missing data did not significantly alter the results. This suggests the results are robust, irrespective of missing data.

Categorical data were analysed using Chi-square test and logistic regression. Continuous data were compared using a T-test, or where data were ordinal, or non-normally distributed, a non-parametric alternative. The level of significance for all tests was set to  $p < 0.05$ . The mean and standard deviation (SD) or standard error (SE) are reported for all normally distributed data and the median for all non-normally distributed data.

Logistic regression was undertaken where appropriate by entering all univariate predictors where  $p < 0.2$  using a backward sequential approach. Odds ratios (OR) and adjusted OR resulting from logistic regression were reported alongside 95% CI (confidence interval). A fully adjusted model is also presented, where univariate predictors with a  $p$  value  $< 0.1$  were included.

It is uncertain if “mild COM” changes (stage one retraction of the pars tensa or flaccida) represent temporary changes in the appearance of the TM, or reflect a milder form of COM. For this clinical based reason, two sets of sub group analyses were undertaken, comparing:

Model 1: No COM vs. mild or significant COM

Model 2: No or mild COM vs. significant COM

## **Results and Analysis**

Full information on the number of ALSPAC participants can be found elsewhere<sup>21, 29</sup>. 14,062 surviving children (at one year) were recruited to the study (Figure 1). Of these, 6,935 (49%) children attended the ALSPAC Focus clinic at age nine (Figure 1). Female children, who were white, with a higher parental social class and higher education level of the mother, were all more likely to attend a clinic visit (all  $p < 0.001$ ).

Twenty-six parents withdrew their children from the study and 19 children had cleft palate. These cases were removed from analysis.

During the clinic visit, COM data (video-otoscopic imaging) could not be obtained for either ear for 330 (5%) children (reasons included child and/or parent declined, equipment availability and time constraints)<sup>20</sup>. No difference was found in any variable between these children and those that did have data for at least one ear. Therefore these 330 children were excluded. There were thus 6560 participants for analysis.

There were 5814 (89%) participants who had no evidence of COM. Of the remaining participants, 350 (5%) were categorised with mild COM and 396 (6%) with significant COM.

Of those with significant COM, 13 participants were categorised with significant COM due to a clinical history of repaired TM perforation; 9 due to a history of cholesteatoma. The remaining were characterised from results of video-otoscopic imaging.

Participant and parent demographics, for both the original data and the imputed plus original data, are presented in Table 1. The ratio of females to males was split evenly (Table 1). The mean age of the mother at birth was 29 (SE 0.07) years old, ranging from 15-44 years old. ENT derived descriptive data, for both the original data and the imputed data, are presented in supplementary Tables A, and additional descriptive data are presented in supplementary Table B.

Table 2 presents the univariate comparisons which showed evidence of a significant association between children with and without COM (any associations that were tested and deemed not to show evidence of significance are presented in the supplementary material). Parental social class, sex of the child and maternal educational level were not associated with the presence of COM. Age of mother at birth was associated with the presence of COM (only when examined as a continuous variable) although the mean difference between outcome groups was small (0.4 and 0.6 years for model 1 and 2 respectively) (Table 2). A number of ENT variables showed increasing evidence of being a significant univariate predictors of COM. These included a history of pus from the child's ear, snoring, previous grommet insertion, adenoidectomy, tonsillectomy, type B or C tympanograms and hearing loss at age 7 (Table 2).

For the majority of additional variables, there was no strong evidence of an association with COM (supplementary material), with the exception of lower birthweight and earlier gestational age at birth. Allergy, asthma, attendance at day-care, crowding, parental smoking, number of siblings and use of dummy/pacifier were not associated with presence of COM (Table 2).

Multivariate models are presented in Table 3. For both models significant independent predictors included an early history of pus/mucus, a history of grommet insertion and a gestational age <37 weeks (Table 3). Type B or C tympanometric recordings at age 7 were also an independent predictor of COM (Table 3).

The fully adjusted model for no COM vs. mild COM/significant COM (all factors with a univariate  $p$  value  $< 0.01$ ) is presented in supplementary table C. It can be seen that the models, and OR's remain very similar to the independent model, implying that the models are stable and robust.

## **Discussion**

To the authors' knowledge, this is the largest prospectively collected cohort study examining a wide range of social and health factors associated with the development of COM in children by the age of nine.

A comparison of results from the two-analysis models demonstrates that the group with "mild COM" have a profile more similar to the "No COM" group. The fact that children with mild COM share more traits with the "no COM" group, suggests that mild retraction of the tympanic membrane may be transient change and that progression to more significant COM is not inevitable. This may have implications for the clinical assessment and management of early TM retraction in the paediatric population and would support a conservative management approach unless more significant signs and symptoms of COM become apparent.

Factors that were univariately associated with the development of COM by the age of 9 included a history of ear discharge, prolonged snoring, grommet insertion (especially multiple insertions), adenoidectomy and/or tonsillectomy, abnormal tympanometry (especially type B), hearing loss at age 7, low birthweight and gestational age under 37 weeks. Multivariate analysis of these factors reveals a more refined picture. Multiple variables show a high degree of multicollinearity (both clinically and statistically). ENT factors of a history of pus/mucus from the ears, previous grommet insertion and type B or C tympanometry remained independent predictors of COM after statistical adjustment. This suggests that the failure of other variables to enter the models (e.g. snoring) is probably due to mediating or confounding relationships between the ENT variables.

Although snoring, adenoidectomy and tonsillectomy are significant univariate predictors of COM, this significance is lost in the multivariate analysis. This appears to be due to the fact that it is highly related to a number of other variables (pus/mucus leakage, breath holding, mother reported child having had grommets, tonsillectomy and adenoidectomy). This interrelationship is not unexpected; the link between adeno-tonsillar hypertrophy and sleep disordered breathing (snoring and obstructive sleep apnoea) in children is well established. Adeno-tonsillectomy remains the primary treatment for paediatric obstructive sleep apnoea<sup>30-32</sup>. A link between COM and adenoidectomy has been proposed by previous studies. A 2010 retrospective cohort study that suggested adenoidectomy may have a protective role in reducing the risk of developing paediatric COM<sup>14</sup>. Our findings suggest that the adenoidectomy, tonsillectomy and nocturnal airway obstruction may have roles to play as risk factors for the development of COM, although the pathophysiology underpinning this unclear. Tonsillectomy and adenoidectomy were also related to whether the child had grommets inserted. This may provide further reason why they were not independent predictors of COM during multivariate analysis. UK studies have shown that adenoidectomy at the time of grommet insertion reduces the risk of recurrent OME<sup>33</sup>.

The association between COM and abnormal tympanometry supports the concepts that middle ear pressure homeostasis and ET function may be important in the formation of COM. The lower odds ratio found in patients with type C1/2 compared to type B tympanograms, and only a single grommet insertion compared to multiple insertions would suggest a dose response relationship between severity of ET dysfunction/OME and risk of COM. The relationship between the physiology of the adenoids, ET and the formation of COM is likely to be complex and cannot be directly quantified in this study. Our findings, in combination with previous studies lend support to an interaction between the three. Various models of COM pathogenesis have been proposed<sup>34-42</sup>. One of the principle theories proposes COM is related to poor middle ear ventilation via Eustachian tube (ET) dysfunction. This theory suggests that chronic lack of middle ear ventilation results in negative middle ear pressures. In combination with multiple infections and weakening of the tympanic membrane, this results in progressive retraction of the ear drum<sup>40</sup>. Entrapment of keratin (produced by the surface of the TM) within the pocket leads to formation of the cholesteatoma. Added infection through microbial colonisation and progressive erosion of middle ear structures results in the typical clinical symptoms of hearing loss and otorrhea.

This retraction theory of cholesteatoma is supported by evidence that cholesteatoma is often associated with ET dysfunction<sup>43, 44</sup>. In patients with COM, a high proportion show COM findings in the contralateral ear<sup>45, 46</sup>. COM, in all of its forms, is particularly present in patients with cleft palate where ET function is known to be disrupted<sup>22, 23</sup>. COM is associated with particular variations in ET anatomy and can be more prevalent in certain ethnic groups<sup>2</sup>.

Whilst this study demonstrates associations between ENT variables and the development of COM by the age of nine, it is not appropriate to suggest a causal relationship. Some of the factors measured may be a result of COM itself, including ear discharge, hearing loss and abnormal tympanometry or audiometry, and as such should be considered as markers of risk of associated COM. The association between COM and some surgical interventions (including grommet insertion, adenoidectomy and tonsillectomy) may also reflect medical treatment of the condition itself.

A strong association was observed between preterm birth and presence of COM; to our knowledge this finding has not previously been reported in the literature although associations between preterm delivery, AOM and OME have been identified<sup>47-49</sup>.

Although age of mother was associated with COM when used as a continuous variable, the mean difference was very small, and it was not considered to be clinically significant. It is probable that the small differences are due to an increased sample size and therefore a high power, rather than any statistical difference. Age of the mother, as a categorical variable, was not an independent predictor of COM (Table 2a).

A number of factors, suggested by previous studies to be associated with the formation of COM, were not supported by our results<sup>6, 8-11, 13, 15-17</sup>. No association was found between COM and sex of the child, parental social class, parental smoking, child-care arrangements, crowding, number of siblings, maternal education level or a history of respiratory infections (see supplementary material). In 2015, Khalid-Raja et al examined UK Hospital Episode

Statistics data for cholesteatoma surgery and found a positive association with the deprivation index of the health authority<sup>50</sup>. No relationship between the presence of COM at the age of 9 and indicators of social deprivation were found here when using the social circumstances of the family rather than of the area.

The strengths of this study are that it is to our knowledge the largest prospective population study on this topic, and as such less prone to biases inherent in smaller retrospective studies. The video-otoscopic images of the TM provide a measure of COM that does not rely on parental report or surgical treatment. Limitations of the study include the fact that children attending the 9 year review clinic (at which otoscopic images were taken) were more likely to be female, white, have a higher parental social class and higher education level of the mother. However these factors were not associated with COM in our analyses; consequently the selection bias is unlikely to have influenced the nature of the associations measured.

Patterns of COM differ with world geography and ethnicity<sup>2</sup>. As such, this study may only identify associations for UK or similar populations. The strong association with multiple grommet insertions, type B/C tympanograms, and ear discharge of pus would seem to support the role of Eustachian tube dysfunction and prolonged OME as a risk factor for developing COM<sup>40, 43-46</sup>.

## **Conclusion**

This is the largest prospective cohort study looking at the associations between the development of paediatric COM (by the age of 9) and health and social factors. The key findings are strong associations between the presence of COM at age 9 and early reported history of otitis media, tympanometric evidence of middle ear effusion and negative middle ear pressure at age 7, hearing loss at age 7 and a history of grommet treatment. Preterm birth and low birthweight were also predictive factors. An association was also found between the development of COM and prolonged snoring, adenoidectomy and/or tonsillectomy.

Some associations suggested by previous published studies (sex of the child, socioeconomic group, parental smoking, maternal education, attendance at childcare, crowding, number of siblings) were not found to be significant predictors in this analysis.

It is not possible through this study method to show a causal relationship between COM and the variables measured. A number of these variables may be related to symptoms or treatment of suspected chronic ear disease. The evidence presented in this study would support a role for prematurity, low birth weight, early acute ear disease, Eustachian tube pathology and upper airway resistance (snoring, adenoidectomy, tonsillectomy) in the pathogenesis of COM.

### **Acknowledgements**

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team (including interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses). This study is based on the original work performed by Mr DD Pothier and Mr AR Maw who reviewed and categorised the otoscopic images for the ALSPAC study.

### **Funding**

The UK Medical Research Council and the Wellcome Trust (**Grant ref: 102215/2/13/2**) and the University of Bristol currently provide core support for ALSPAC. Funding for the analyses in this paper were generously provided through a research grant from the “Above and Beyond” charity for central Bristol Hospitals.

The authors have no competing interests to declare.

### **Summary**

- It is unclear why some children develop chronic otitis media (COM).



- 6560 children underwent otoscopic photography at the age 9 as part of a prospective, longitudinal cohort study.
- 5% of children were diagnosed with mild COM and 6% with more significant COM
- The development of COM by the age of 9 was associated with a history of otorrhea, snoring, grommet insertion, adenoidectomy, tonsillectomy, hearing loss, abnormal tympanograms and preterm birth.
- Some associations suggested by previous published studies (sex of the child, socioeconomic group, parental smoking, maternal education, attendance at childcare, crowding, number of siblings) were not found to be significant predictors in this analysis.
- This represents the largest prospective cohort study of factors associated with the development of paediatric chronic otitis media to date.

## **References**

1. Teele DW, Klein JO, Rosner B. Epidemiology of otitis media during the first seven years of life in children in greater Boston: a prospective, cohort study. *J Infect Dis* 1989;**160**:83-94
2. Bluestone CD. Epidemiology and pathogenesis of chronic suppurative otitis media: implications for prevention and treatment. *Int J Pediatr Otorhinolaryngol* 1998;**42**:207-23
3. Monasta L, Ronfani L, Marchetti F, Montico M, Vecchi Brumatti L, Bavcar A, et al. Burden of disease caused by otitis media: systematic review and global estimates. *PLoS One* 2012;**7**:e36226
4. Acuin J, World Health Organisation. Chronic suppurative otitis media : Burden of illness and management options. *World Health Organisation, Geneva, Switzerland* 2004
5. Nunez DA, Browning GG. Risks of developing an otogenic intracranial abscess. *J Laryngol Otol* 1990;**104**:468-72
6. Browning GG, Gatehouse S. The prevalence of middle ear disease in the adult British population. *Clin Otolaryngol Allied Sci* 1992;**17**:317-21
7. Elemraid MA, Mackenzie IJ, Fraser WD, Harper G, Faragher B, Atef Z, et al. A case-control study of nutritional factors associated with chronic suppurative otitis media in Yemeni children. *Eur J Clin Nutr* 2011;**65**:895-902
8. Fliss DM, Shoham I, Leiberman A, Dagan R. Chronic suppurative otitis media without cholesteatoma in children in southern Israel: incidence and risk factors. *Pediatr Infect Dis J* 1991;**10**:895-9
9. Koch A, Homoe P, Pipper C, Hjuler T, Melbye M. Chronic suppurative otitis media in a birth cohort of children in Greenland: population-based study of incidence and risk factors. *Pediatr Infect Dis J* 2011;**30**:25-9
10. Lasisi AO, Olaniyan FA, Muibi SA, Azeez IA, Abdulwasiiu KG, Lasisi TJ, et al. Clinical and demographic risk factors associated with chronic suppurative otitis media. *Int J Pediatr Otorhinolaryngol* 2007;**71**:1549-54
11. Park M, Lee JS, Lee JH, Oh SH, Park MK. Prevalence and risk factors of chronic otitis media: the Korean National Health and Nutrition Examination Survey 2010-2012. *PLoS One* 2015;**10**:e0125905
12. Santos-Cortez RL, Reyes-Quintos MR, Tantoco ML, Abbe I, Llanes EG, Ajami NJ, et al. Genetic and Environmental Determinants of Otitis Media in an Indigenous Filipino Population. *Otolaryngol Head Neck Surg* 2016;**155**:856-62

13. Shaheen MM, Raquib A, Ahmad SM. Prevalence and associated socio-demographic factors of chronic suppurative otitis media among rural primary school children of Bangladesh. *Int J Pediatr Otorhinolaryngol* 2012;**76**:1201-4
14. Spilsbury K, Miller I, Semmens JB, Lannigan FJ. Factors associated with developing cholesteatoma: a study of 45,980 children with middle ear disease. *Laryngoscope* 2010;**120**:625-30
15. Wang J, Chen B, Xu M, Wu J, Wang T, Zhao J, et al. Etiological factors associated with chronic suppurative otitis media in a population of Han adults in China. *Acta Otolaryngol* 2016;**136**:1024-8
16. Zakzouk SM, Hajjaj MF. Epidemiology of chronic suppurative otitis media among Saudi children--a comparative study of two decades. *Int J Pediatr Otorhinolaryngol* 2002;**62**:215-8
17. van der Veen EL, Schilder AG, van Heerbeek N, Verhoeff M, Zielhuis GA, Rovers MM. Predictors of chronic suppurative otitis media in children. *Arch Otolaryngol Head Neck Surg* 2006;**132**:1115-8
18. Taipale A, Pelkonen T, Taipale M, Bernardino L, Peltola H, Pitkaranta A. Chronic suppurative otitis media in children of Luanda, Angola. *Acta Paediatr* 2011;**100**:e84-8
19. Zhang Y, Xu M, Zhang J, Zeng L, Wang Y, Zheng QY. Risk factors for chronic and recurrent otitis media-a meta-analysis. *PLoS One* 2014;**9**:e86397
20. Maw AR, Hall AJ, Pothier DD, Gregory SP, Steer CD. The prevalence of tympanic membrane and related middle ear pathology in children: a large longitudinal cohort study followed from birth to age ten. *Otol Neurotol* 2011;**32**:1256-61
21. Boyd A, Golding J, Macleod J, Lawlor DA, Fraser A, Henderson J, et al. Cohort Profile: the 'children of the 90s'--the index offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol* 2013;**42**:111-27
22. Djurhuus BD, Skytthe A, Faber CE, Christensen K. Cholesteatoma risk in 8,593 orofacial cleft cases and 6,989 siblings: A nationwide study. *Laryngoscope* 2015;**125**:1225-9
23. Sheahan P, Blayney AW, Sheahan JN, Earley MJ. Sequelae of otitis media with effusion among children with cleft lip and/or cleft palate. *Clin Otolaryngol Allied Sci* 2002;**27**:494-500
24. Browning GG, S.N. M, Kelly G, Swan IRC, Canter R, Mckerrow WS. 237c: Chronic otitis media. In: Gleeson M, ed. *Scott-Brown's Otorhinolaryngology, Head and Neck Surgery (7th edition)*: Hodder Arnold, London, UK. , 2008;3395-445
25. Roberts G, Peckitt C, Northstone K, Strachan D, Lack G, Henderson J, et al. Relationship between aeroallergen and food allergen sensitization in childhood. *Clin Exp Allergy* 2005;**35**:933-40

26. Williams HC, Burney PG, Hay RJ, Archer CB, Shipley MJ, Hunter JJ, et al. The U.K. Working Party's Diagnostic Criteria for Atopic Dermatitis. I. Derivation of a minimum set of discriminators for atopic dermatitis. *Br J Dermatol* 1994;**131**:383-96
27. Williams HC, Burney PG, Pembroke AC, Hay RJ. The U.K. Working Party's Diagnostic Criteria for Atopic Dermatitis. III. Independent hospital validation. *Br J Dermatol* 1994;**131**:406-16
28. Williams HC, Burney PG, Strachan D, Hay RJ. The U.K. Working Party's Diagnostic Criteria for Atopic Dermatitis. II. Observer variation of clinical diagnosis and signs of atopic dermatitis. *Br J Dermatol* 1994;**131**:397-405
29. Fraser A, Macdonald-Wallis C, Tilling K, Boyd A, Golding J, Davey Smith G, et al. Cohort Profile: the Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *Int J Epidemiol* 2013;**42**:97-110
30. Brockbank JC. Update on pathophysiology and treatment of childhood obstructive sleep apnea syndrome. *Paediatr Respir Rev* 2017
31. Chinnadurai S, Jordan AK, Sathe NA, Fonnesbeck C, McPheeters ML, Francis DO. Tonsillectomy for Obstructive Sleep-Disordered Breathing: A Meta-analysis. *Pediatrics* 2017;**139**
32. Ehsan Z, Ishman SL. Pediatric Obstructive Sleep Apnea. *Otolaryngol Clin North Am* 2016;**49**:1449-64
33. Group MRCMOMS. Adjuvant adenoidectomy in persistent bilateral otitis media with effusion: hearing and revision surgery outcomes through 2 years in the TARGET randomised trial. *Clin Otolaryngol* 2012;**37**:107-16
34. Persaud R, Hajioff D, Trinidad A, Khemani S, Bhattacharyya MN, Papadimitriou N, et al. Evidence-based review of aetiopathogenic theories of congenital and acquired cholesteatoma. *J Laryngol Otol* 2007;**121**:1013-9
35. Hamed MA, Nakata S, Sayed RH, Ueda H, Badawy BS, Nishimura Y, et al. Pathogenesis and Bone Resorption in Acquired Cholesteatoma: Current Knowledge and Future Prospectives. *Clin Exp Otorhinolaryngol* 2016;**9**:298-308
36. Jackler RK, Santa Maria PL, Varsak YK, Nguyen A, Blevins NH. A new theory on the pathogenesis of acquired cholesteatoma: Mucosal traction. *Laryngoscope* 2015;**125 Suppl 4**:S1-S14
37. Jennings BA, Prinsley P, Philpott C, Willis G, Bhutta MF. The genetics of cholesteatoma. A systematic review using narrative synthesis. *Clin Otolaryngol* 2017

38. Louw L. Acquired cholesteatoma pathogenesis: stepwise explanations. *J Laryngol Otol* 2010;**124**:587-93
39. Olszewska E, Wagner M, Bernal-Sprekelsen M, Ebmeyer J, Dazert S, Hildmann H, et al. Etiopathogenesis of cholesteatoma. *Eur Arch Otorhinolaryngol* 2004;**261**:6-24
40. Ramakrishnan Y, Kotecha A, Bowdler DA. A review of retraction pockets: past, present and future management. *J Laryngol Otol* 2007;**121**:521-5
41. Wells MD, Michaels L. Role of retraction pockets in cholesteatoma formation. *Clin Otolaryngol Allied Sci* 1983;**8**:39-45
42. Yoon TH, Schachern PA, Paparella MM, Aeppli DM. Pathology and pathogenesis of tympanic membrane retraction. *Am J Otolaryngol* 1990;**11**:10-7
43. Bluestone CD, Cantekin EI, Beery QC, Stool SE. Function of the Eustachian tube related to surgical management of acquired aural cholesteatoma in children. *Laryngoscope* 1978;**88**:1155-64
44. Yuceturk AV, Unlu HH, Okumus M, Yildiz T, Filiz U. The evaluation of eustachian tube function in patients with chronic otitis media. *Clin Otolaryngol Allied Sci* 1997;**22**:449-52
45. Hassman-Poznanska E, Kurzyrna A, Trzpis K, Poznanska M. The status of the contralateral ear in children with acquired cholesteatoma. *Acta Otolaryngol* 2012;**132**:404-8
46. Jadia S, Mourya A, Jain L, Qureshi S. Pre and Post Operative Status of Contralateral Ear in Unilateral Chronic Otitis Media. *Indian J Otolaryngol Head Neck Surg* 2016;**68**:20-4
47. Engel J, Anteunis L, Volovics A, Hendriks J, Marres E. Prevalence rates of otitis media with effusion from 0 to 2 years of age: healthy-born versus high-risk-born infants. *Int J Pediatr Otorhinolaryngol* 1999;**47**:243-51
48. Engel JA, Straetmans M, Zielhuis GA. Birth characteristics and recurrent otitis media with effusion in young children. *Int J Pediatr Otorhinolaryngol* 2005;**69**:533-40
49. Korvel-Hanquist A, Koch A, Niclasen J, Dammeyer J, Lous J, Olsen SF, et al. Risk Factors of Early Otitis Media in the Danish National Birth Cohort. *PLoS One* 2016;**11**:e0166465
50. Khalid-Raja M, Tikka T, Coulson C. Cholesteatoma: a disease of the poor (socially deprived)? *Eur Arch Otorhinolaryngol* 2015;**272**:2799-805

### **Legends for Figures and Tables**

Figure 1. Study flow chart

Table 1. Demographics data of the study sample (n=6560)

Table 2a. Univariate comparisons of demographics data with the outcome variable (original plus imputed data n= 6560)

Table 2b. Univariate comparison of ENT questionnaire data with the outcome variable (original plus imputed data n= 6560)

Table 2c. Comparison of additional variables with the outcome variables (original plus imputed data n= 6560)


Table 3. Multivariate models (original plus imputed data n= 6560)

Supplementary Table A. ENT derived data(n=6560)

Supplementary Table B. Additional variables (n=6560)

Supplementary Table C. Fully adjusted model (original plus imputed data n= 6560).

**Figure 1. Study flow chart**



\*percentage of those who attended a clinic visit.

\*\*There were no significant differences in demographic data between those participants who had ear data and those that did not: Gender ( $p = 0.827$ ), Ethnic background ( $p = 0.655$ ), social class ( $p = 0.942$ ) and mother's highest education level ( $p = 0.759$ ).

**Table 1. Demographics data of the study sample (n=6560)**

		Original data n (%)*	Original plus imputed data n (%)*
Sex of child	Male	3251 (50)	3251 (50)
	Female	3309 (50)	3309 (50)
	Missing	0	
Child ethnic background	White	5655 (96)	6281 (96)
	Non-white	230 (4)	279 (4)
	Missing	675	
Parental social class <sup>1</sup> (Social class of the most advantaged parent)	I (professional)	897 (16)	958 (15)
	II	2597 (45)	2861 (44)
	III (non-manual)	1388 (24)	1598 (24)
	III (manual)	581 (10)	717 (11)
	IV (partly skilled)	220 (4)	277 (4)
	V (unskilled)	31 (<1)	149 (2)
	Missing	846	
Mothers highest education qualification	CSE (lowest) <sup>^</sup>	605 (11)	779 (12)
	Vocational <sup>^</sup>	515 (9)	621 (9)
	O level (medium) <sup>^^</sup>	2068 (36)	2357 (36)
	A level <sup>^^^</sup>	1618 (28)	1752 (27)
	Degree (highest) <sup>^^^</sup>	970 (17)	1051 (16)
	Missing	784	
Age of mother at birth (years)	Mean	29 (SD 4.6)	29 (SE 0.07)
	Range	15-44	15-44
	Missing	349	

\*unless otherwise stated

<sup>^</sup> Education finished <16 years. <sup>^^</sup> Education up to 16 years. <sup>^^^</sup> Education at ≥18 years

SD - Standard deviation

SE – Standard error



<sup>1</sup> The parents' social class was based on the Standard Occupational Classification (1990) of the mother or partner (whichever was the more advantaged) occupation at 32 weeks of pregnancy<sup>29</sup>. This was measured using the parents' occupation and based on a scale of I (professional), through II, III (divided into manual and non-manual), IV (partly-skilled occupations) and V (unskilled occupations).

Table 2a. Univariate comparisons of demographics data with the outcome variable (original plus imputed data n= 6560)

		n (%) <sup>*</sup>			P value ( <sup>*</sup> )	
		No COM (n=5814)	Mild COM (n=350)	Significant COM (n=396)	No vs. mild / significant	No / mild vs. significant
Sex of child	Male	2894 (50)	171 (49)	186 (47)	0.323 (OR 1.1 [0.9-1.3])	0.288 (OR 1.1 [0.9-1.4])
	Female	2920 (50)	179 (51)	210 (53)		
Child ethnic background	White	5563 (96)	335 (96)	383 (97)	0.535 (OR 0.8 [0.5-1.5])	0.409 (OR 0.7 [0.4-1.5])
	Non-white	251 (4)	15 (4)	13 (3)		
Parental social class (highest parental class of the most advantaged parent)	I (professional)	856 (15)	48 (14)	54 (14)	0.357	0.692
	II	2529 (43)	148 (42)	184 (46)		
	III (non-manual)	1440 (25)	82 (23)	76 (19)		
	III (manual)	625 (11)	41 (12)	51 (13)		
	IV (partly skilled)	241 (4)	20 (6)	16 (4)		
	V (unskilled)	123 (2)	11 (3)	15 (4)		
Mothers highest education qualification	CSE/Vocational (low) <sup>^</sup>	1235 (22)	79 (14)	86 (13)	0.440	0.264
	O level (medium) <sup>^^</sup>	2105 (36)	125 (36)	128 (32)		
	A level/ Degree (high) <sup>^^^</sup>	2474 (27)	146 (25)	182 (31)		
Age of mother at birth (years)	Mean (SE)	29.0 (0.07)	28.9 (0.26)	28.4 (0.22)	<b>0.024</b> <b>(MD 0.4 [0.1-0.8])</b>	<b>0.010</b> <b>(MD 0.6 [0.1-1.1])</b>
Age of mother at birth	≤ 25 years	915 (16)	57 (16)	59 (15)	0.194	0.071
	25 – 34 years	4190 (72)	252 (72)	303 (77)		
	≥ 35 years	709 (12)	41 (12)	34 (9)		

<sup>\*</sup>OR (odds ratio) with 95% CI (confidence intervals) or MD (mean difference) with 95% CI presented, as appropriate. If not significant, P values only, resulting from the Chi-squared test, are reported for tables greater than 2x2.

COM = chronic otitis media

#### Significant result

<sup>^</sup> Education finished <16 years. <sup>^^</sup> Education up to 16 years. <sup>^^^</sup> Education at ≥18 years

Table 2b. Univariate comparison of ENT questionnaire data with the outcome variable (original plus imputed data n= 6560)

		n (%)*			P value (OR [95% CI])*	
Reported**		No COM (n=5814)	Mild COM (n=350)	Significant COM (n=396)	No vs. mild / significant	No / mild vs. significant
Pus/mucus leaking from the child's ear	Never <sup>R</sup>	4439 (76)	207 (59)	197 (50)		
	Once	863 (14)	66 (19)	88 (22)	<b>&lt; 0.001 (2.0 [1.6-2.4])</b>	<b>&lt; 0.001 (2.2 [1.7-2.9])</b>
	> once	512 (9)	77 (22)	111 (28)	<b>&lt; 0.001 (4.0 [3.3-4.9])</b>	<b>&lt; 0.001 (4.4 [3.5-5.7])</b>
Child holds breathe for several seconds while sleeping <sup>1</sup>	No	4070 (70)	241 (69)	271 (68)		
	Yes	1744 (30)	109 (31)	125 (32)	0.478 (1.1 [0.9-1.3])	0.533 (1.1 [0.8-1.6])
Child ever snores <sup>1</sup>	No	4887 (84)	275 (79)	287 (72)		
	Yes	927 (16)	75 (21)	109 (28)	<b>&lt; 0.001 (1.7 [1.4-2.1])</b>	<b>&lt; 0.001 (2.0 [1.5-2.5])</b>
Child snores on at least two time points (prolonged snorting) <sup>2</sup>	No	5277 (91)	302 (86)	323 (82)		
	Yes	537 (9)	48 (14)	73 (18)	<b>&lt; 0.001 (1.9 [1.7-2.1])</b>	<b>&lt; 0.001 (2.1 [1.6-2.8])</b>
Child snores at 6 and 18 months (early snoring) <sup>1</sup>	No	5347 (92)	321 (92)	355 (90)		
	Yes	467 (8)	29 (8)	41 (10)	0.286 (1.2 [0.9-1.6])	0.132 (1.3 [0.9-1.9])
Mother reported a history of having grommets inserted	No	5701 (98)	341 (97)	382 (96)		
	Yes	113 (2)	9 (3)	14 (4)	0.144 (1.6 [0.8-3.0])	0.097 (1.8 [0.9-3.7])
Child had grommets inserted (by age 9)	Did not have grommets <sup>R</sup>	5634 (97)	275 (79)	288 (73)		
	Grommets inserted once	150 (3)	65 (19)	78 (20)	<b>&lt; 0.001 (9.6 [7.5-12.2])</b>	<b>&lt; 0.001 (7.4 [5.6-9.9])</b>
	Grommets inserted >1	30 (<1)	10 (3)	30 (8)	<b>&lt; 0.001 (13.4 [8.3-21.9])</b>	<b>&lt; 0.001 (15.5 [9.5-25.3])</b>
Child had tonsils removed (by age 9)	No	5738 (99)	332 (95)	375 (95)		
	Yes	76 (1)	18 (5)	21 (5)	<b>&lt; 0.001 (4.1 [2.8-6.2])</b>	<b>&lt; 0.001 (3.6 [2.2-5.9])</b>
Child had adenoids removed (by age 9)	No	5692 (98)	315 (90)	345 (87)		
	Yes	122 (2)	35 (10)	51 (13)	<b>&lt; 0.001 (6.1 [4.6-8.1])</b>	<b>&lt; 0.001 (5.6 [4.1-7.9])</b>
Cough count <sup>3</sup>	< 50%	1764 (30)	89 (25)	123 (31)		
	≥ 50%	4050 (70)	261 (75)	273 (69)	0.285 (1.1 [0.9-1.3])	0.682 (0.9 [0.8-1.2])

Fever reported alongside a cough	No cough reported / never	1476 (25)	79 (23)	110 (28)	0.970	0.382
	Fever with some attacks	3263 (56)	197 (56)	222 (56)		
	Fever with every attack	1075 (18)	74 (21)	64 (16)		
Tympanometry at age 7 (better functioning ear)	Type A <sup>R</sup>	4464 (77)	185 (53)	158 (40)	< 0.001 (3.1 [2.4-4.1])	< 0.001 (3.5 [5.6-4.8])
	Type C1 / C2	1037 (18)	101 (29)	118 (30)		
	Type B	313 (5)	64 (18)	119 (30)		
Audiometry age 7 (better hearing ear), dB	< 20 dB	5694 (98)	333 (95)	363 (92)	< 0.001 (3.4 [2.3-5.1])	< 0.001 (4.0 [2.6-6.2])
	≥ 20 dB	120 (2)	17 (5)	33 (8)		

\* OR (odds ratio) with 95% CI (confidence intervals) presented as appropriate. If not significant, P values only, resulting from the Chi-squared test, are reported for tables greater than 2x2.

\*\* All variables are derived from longitudinal questionnaire data reported by the parent(s) when the child was 6, 18, 30, 42, 57, 69, 81, 91 and 103 months.

<sup>1</sup> Yes = a report of yes for at least one minute at least one 'yes often' / 'most nights' across all time points.

<sup>2</sup> Yes = a report of snoring for at least one minute at least two reports of 'most nights' across time points.

<sup>3</sup> Cough count = % of times a cough was reported across all time points.

<sup>R</sup> Reference category

### Significant result

COM= chronic otitis media

Table 2c. Comparison of additional variables with the outcome variables (original plus imputed data n= 6560)

		n (%)*			P value (OR* [95% CI])	
<b>Reported**</b>		No COM (n=5814)	Mild COM (n=350)	Significant COM (n=396)	No vs. mild / significant	No / mild vs. significant
<b>Medical factors</b>						
Allergy <sup>1</sup> (7 years)	No	4575 (79)	282 (81)	319 (81)	0.264 (0.9 [0.7-1.1])	0.384 (0.9 [0.7-1.2])
	Yes	1239 (21)	68 (19)	77 (19)		
Atopic dermatitis (7 years)	No	5347 (92)	320 (91)	366 (92)	0.981 (1.0 [0.7-1.3])	0.786 (0.9 [0.6-1.4])
	Yes	467 (8)	30 (9)	30 (8)		
Doctor diagnosis of asthma (91 months)	No	4617 (79)	278 (79)	310 (78)	0.703 (1.0 [0.9-1.3])	0.594 (1.1 [0.8-1.4])
	Yes	1197 (21)	72 (21)	86 (22)		
<b>Day care</b>						
Day care at any time (up to 38 months) <sup>2</sup>	None	3498 (60)	203 (58)	235 (59)	0.525 (1.1 [0.9-1.3])	0.853 (1.0 [0.8-1.3])
	Yes	2316 (40)	147 (42)	160 (41)		
Day care regularly (38 months) <sup>3</sup>	No	3862 (66)	222 (63)	261 (66)	0.317 (1.1 [0.9-1.3])	0.685 (1.1 [0.8-1.3])
	Yes	1952 (34)	128 (37)	135 (34)		
Day care 10+ hours per week (38 months)	No	1051 (18)	67 (19)	64 (16)	0.671 (1.0 [0.8-1.3])	0.362 (1.2 [0.8-1.7])
	Yes	4763 (82)	283 (81)	332 (84)		
<b>Crowding</b>						
Number of other children in house (6 months)	0	2613 (45)	133 (38)	180 (45)	0.261	0.848
	1	2153 (37)	153 (44)	148 (37)		
	>2	1047 (18)	65 (19)	68 (17)		
Number of other children in house (81 months)	0	606 (10)	31 (9)	36 (9)	0.430	0.413
	1	3137 (54)	185 (53)	225 (57)		
	>2	2071 (36)	134 (38)	136 (34)		

Crowding index (8 months) <sup>4</sup>	≤0.5	911 (16)	56 (14)	54 (16)	0.264	0.720
	0.5 - 0.75	2223 (37)	121 (36)	135 (37)		
	0.75 - 1	1890 (32)	121 (33)	150 (30)		
	>1	788 (15)	52 (17)	58 (17)		
Crowding index (33 months) <sup>4</sup>	≤0.5	1127 (19)	67 (19)	75 (19)	0.747	0.724
	0.5 - 0.75	2185 (38)	136 (39)	153 (39)		
	0.75 - 1	2107 (36)	123 (35)	140 (35)		
	>1	396 (7)	23 (7)	29 (7)		
Crowding index (81 months) <sup>4</sup>	≤0.5	778 (14)	57 (14)	52 (15)	0.720	0.382
	0.5 - 0.75	1962 (31)	117 (31)	128 (31)		
	0.75 - 1	2293 (35)	128 (38)	157 (35)		
	>1	781 (19)	48 (18)	60 (20)		
Presence of mould in the home (8 months)	None <sup>R</sup>	4069 (70)	228 (65)	267 (67)	0.101	0.366
	Not serious	1274 (22)	93 (27)	90 (23)		
	Serious	471 (8)	29 (8)	39 (10)		
Presence of damp in the home (8 months)	None <sup>R</sup>	2836 (49)	168 (48)	192 (48)	0.818	0.851
	Not serious	2585 (44)	157 (45)	175 (44)		
	Serious	393 (7)	24 (7)	29 (7)		
<b>Smoking</b>						
Child exposure to smoke (up to 54 months) <sup>5</sup>	None	3237 (56)	191 (54)	212 (54)	0.424 (1.1 [0.9-1.3])	0.430 (1.1 [0.9-1.3])
	Yes	2577 (44)	159 (45)	184 (46)		
Child exposure to smoke (6 months)	None	3843 (66)	233 (67)	255 (64)	0.756 (1.0 [0.9-1.2])	0.526 (1.1 [0.9-1.4])
	Yes	1971 (34)	117 (33)	141 (36)		
Child exposure to smoke (38 months)	None	3740 (64)	215 (61)	254 (64)	0.484 (1.1 [0.9-1.3])	0.971 (1.0 [0.8-1.3])
	Yes	2074 (36)	135 (39)	142 (36)		
Child exposure to smoke (54 months)	None	3836 (66)	221 (63)	250 (63)	0.126 (1.1 [1.0-1.3])	0.300 (1.1 [0.9-1.4])
	Yes	1978 (34)	129 (37)	146 (37)		

Maternal smoking during pregnancy (at any time) <sup>6</sup>	None	4700 (77)	284 (78)	326 (78)		
	Yes	1114 (23)	66 (22)	70 (22)	0.584 (0.9 [0.8-1.2])	0.526 (0.9 [0.7-1.2])
Maternal smoking during pregnancy (up to 18 weeks)	None	4604 (79)	276 (79)	318 (80)		
	Yes	1210 (21)	74 (21)	78 (20)	0.775 (1.0 [0.8-1.2])	0.580 (0.9 [0.7-1.2])
Maternal smoking during pregnancy (up to 32 weeks)	None	4620 (79)	275 (79)	318 (80)		
	Yes	1194 (21)	75 (21)	78 (20)	0.991 (1.0 [0.8-1.2])	0.681 (0.9 [0.7-1.2])
<b>At birth</b>						
Breast feeding at any time point <sup>^</sup>	No	991 (17)	64 (18)	74 (19)		
	Yes	4823 (83)	286 (82)	322 (82)	0.387 (0.9 [0.7-1.1])	0.486 (0.9 [0.7-1.2])
Breast feeding duration across time points <sup>^</sup>	Never	1254 (22)	74 (21)	92 (23)		
	< 3 months	1373 (24)	76 (22)	93 (23)		
	3-5 months	997 (17)	59 (17)	68 (17)	0.859	0.474
	> 6 months	2190 (38)	142 (40)	144 (36)		
Birthweight of child (g)	Mean (SE)	3415 (7.8)	3394 (35.0)	3332 (31.1)	<b>0.023</b> <b>(MD 53 [7.6-99.2])</b>	<b>0.009</b> <b>(MD 81.2 [20.3-142.0])</b>
Birthweight of child	> 2500g	5524 (95)	326 (93)	365 (92)	<b>0.015</b>	<b>0.030</b>
	≤ 2500g	290 (5)	24 (7)	31 (8)	<b>(1.5 [1.1-2.1])</b>	<b>(1.6 [1.1-2.4])</b>
Gestational age of child (weeks)	Mean (SE)	39.5 (0.03)	39.3 (0.2)	39.3 (0.1)	<b>0.040</b> <b>(MD 0.2 [0.1-0.3])</b>	0.134 (MD 0.1 [0.1-1.2])
Gestational age of child	≥ 37 weeks	5514 (95)	328 (94)	359 (91)	<b>0.005</b>	<b>0.001</b>
	< 37 weeks	300 (5)	22 (6)	37 (9)	<b>(1.6 [1.1-2.2])</b>	<b>(1.8 [1.3-2.7])</b>
Child ever use a dummy/ pacifier	No	3080 (52)	186 (53)	207 (52)		
	Yes	2734 (47)	164 (47)	188 (48)	0.936 (1.0 [0.9-1.2])	0.850 (1.0 [0.8-1.3])

\* Unless otherwise stated. MD – Mean difference. If not significant, P values only, resulting from the Chi-squared test, are reported for tables greater than 2x2.

\*\* All variables are derived from questionnaire data reported by the parent(s) when the child was 6, 18, 30, 42, 57, 69, 81, 91 and 103 months.

<sup>1</sup> A child was deemed to react to an allergen if the weal and/or flare was  $\geq 2$ mm at the age 7 Focus clinic. Children were excluded if they didn't react to the positive allergen, or did react to the negative control. These excluded children were imputed along with the missing data.

<sup>2</sup> The child attended day care, as reported by the parent, at 15, 24 and 38 months.

<sup>3</sup> 'Regular' was defined as general habit and not a one off.

<sup>4</sup> A measure of the number of persons in the household divided by the number of rooms available (including the kitchen if large enough to eat in).

<sup>5</sup> Child had some exposure to smoking (passive smoking) at any point up to 54 months

<sup>6</sup> Mother reported smoking during pregnancy (at any time).

<sup>^</sup> Owing to a difference in time points, numbers may differ.



Table 3. Multivariate models (original plus imputed data n= 6560)

		n (%)		No COM vs. mild / significant COM	
Model 1		No COM (n=5814)	Mild/significant COM (n=746)	Adjusted <i>p</i> value	Adjusted OR (95% CI)
Tympanometry (age 7)	Type A <sup>R</sup>	4464 (77)	344 (46)	< 0.001	2.4 (1.9-2.9)
	Type C1 / C2	1037 (18)	220 (29)		
	Type B	313 (5)	183 (25)		
Pus/mucus leaking from the child's ear	Never	4439 (76)	404 (54)	< 0.001	1.4 (1.3-1.6)
	At least once	1375 (24)	342 (46)		
Child had grommets inserted (by age 9)	No	5634 (97)	563 (75)	< 0.001	5.0 (3.9-6.5)
	Yes	180 (3)	183 (25)		
Gestational age	≥ 37 weeks	5514 (95)	687 (92)	0.040	1.4 (1.1-1.9)
	< 37 weeks	300 (5)	59 (8)		
Model 2		n (%)		No / mild COM vs. significant COM	
		No/mild COM (n=6164)	Significant COM (n=396)	Adjusted <i>p</i> value	Adjusted OR (95% CI)
Tympanometry (age 7)	Type A <sup>R</sup>	4464 (77)	344 (46)	< 0.001	2.5 (2.0-3.3)
	Type C1 / C2	1037 (18)	220 (29)		
	Type B	313 (5)	183 (25)		
Pus/mucus leaking from the child's ear	Never	4645 (75)	197 (50)	< 0.001	1.9 (1.5-2.3)
	At least once	1519 (25)	199 (50)		
Child had grommets inserted (by age 9)	No	5909 (96)	288 (73)	< 0.001	3.9 (2.9-5.1)
	Yes	255 (4)	108 (27)		
Gestational age	≥ 37 weeks	5842 (95)	359 (91)	0.014	1.6 (1.1-2.4)
	< 37 weeks	322 (5)	37 (9)		

<sup>R</sup> Reference category

COM = chronic otitis media. OR = odds ratio. CI = confidence interval.

Supplementary table A. ENT derived data(n=6560)

<b>Reported**</b>		<b>Original data n (%)*</b>	<b>Original plus imputed data n (%)*</b>
Pus/mucus leaking from the child's ear	Never	4690 (74)	4842 (74)
	Once	998 (16)	1018 (16)
	>1	673 (11)	700 (11)
	Missing	199	
Child holds breathe for several seconds while sleeping <sup>1</sup>	No	4140 (70)	4583 (70)
	Yes	1805 (30)	1977 (30)
	Missing	615	
Child ever snores <sup>1</sup>	No	5274 (83)	5449 (83)
	Yes	1051 (17)	1111 (17)
	Missing	235	
Child snores on at least two time points (prolonged snoring) <sup>2</sup>	No	5807 (92)	5903 (90)
	Yes	518 (8)	657 (10)
	Missing	235	
Child snores at 6 and 18 months (early snoring) <sup>3</sup>	No	5642 (96)	6023 (92)
	Yes	261 (4)	537 (6)
	Missing	657	
Mother reported a history of having grommets inserted	No	5133 (98)	6424 (98)
	Yes	85 (2)	136 (2)
	Missing	1342	
Child had grommets inserted (by age 9)	Did not have grommets	6166 (95)	6197 (94)
	Grommets inserted once	291 (5)	293 (4)
	Grommets inserted >1	66 (1)	70 (1)
	Missing	37	
Child had tonsils removed (by age 9)	No	6414 (98)	6445 (98)
	Yes	112 (2)	115 (2)

	Missing	34	
Child had adenoids removed (by age 9)	No	6322 (97)	6352 (97)
	Yes	204 (3)	208 (3)
	Missing	34	
Cough count <sup>4</sup>	Mean	61 (SD 28)	60 (SE 0.4)
Fever reported alongside a cough <sup>5</sup>	No cough reported / never	1515 (25)	1664 (25)
	Fever with some attacks	3462 (57)	3681 (56)
	Fever with every attack	1155 (19)	1215 (19)
	Missing	428	
Tympanometry at age 7 (better functioning ear) <sup>6</sup>	Type A	4059 (75)	4808 (73)
	Type C1	646 (12)	790 (12)
	Type C2	360 (7)	466 (7)
	Type B	369 (7)	496 (8)
	Missing	1126	
Audiometry age 7 (better hearing ear) <sup>7</sup>	< 20 dB	5100 (98)	6390 (97)
	≥ 20 dB	126 (2)	170 (3)
	Missing	1334	

\*Unless otherwise stated.

\*\* All variables are derived from longitudinal questionnaire data reported by the parent(s) when the child was 6, 18, 30, 42, 57, 69, 81, 91 and 103 months.

<sup>1</sup> Yes = a report of yes for at least one minute, with at least one 'yes often' or 'most nights' across all time points.

<sup>2</sup> Yes = a report of snoring for at least one minute, with at least two reports of 'most nights' across time points.

<sup>3</sup> Categorised as positive if there was a report of snoring 'most nights' at either 6 or 18 months

<sup>4</sup> Cough count = % of times a cough was reported across all time points (cough count was considered missing if > 50% of time points were missing).

<sup>5</sup> Categorised as 'no cough reported or no fever recorded with any cough attack', 'fever with some cough attacks' or 'fever with every cough attack'.

<sup>6</sup> Tympanograms were coded according to Fieullau-Nikolajsen's modification of Jerger's classification, categorised as type A, C1/C2 or B in the better functioning ear<sup>28</sup>.

<sup>7</sup> Hearing thresholds were measured using pure tone audiometry across 0.5, 1, 2 and 4 kHz in the left and right ears. The air conduction average hearing threshold in the better hearing ear was calculated.

SD - Standard deviation, SE – Standard error

Supplementary Table B. Additional variables (n=6560)

Reported**		Original data n (%)*	Original plus imputed data n (%)*
<b>Medical factors</b>			
Allergy (7 years) <sup>1</sup>	No	3875 (79)	5176 (79)
	Yes	1032 (21)	1384 (21)
	Excluded	111	
	Missing	1542	
Atopic dermatitis (7 years) <sup>2</sup>	No	5268 (92)	6032 (92)
	Yes	464 (8)	528 (8)
	Missing	828	
Doctor diagnosis of asthma (91 months)	No	4153 (80)	5205 (79)
	Yes	1050 (20)	1355 (21)
	Missing	1357	
<b>Day care</b>			
Day care at any time (up to 38 months) <sup>3</sup>	None	3021 (57)	3936 (60)
	Yes	2303 (43)	2624 (40)
	Missing	1236	
Day care regularly (38 months) <sup>4</sup>	No	3490 (62)	4100 (63)
	Yes	2112 (38)	2460 (38)
	Missing	958	
Day care 10+ hours per week (38 months)	No	4665 (83)	5378 (82)
	Yes	937 (17)	1182 (18)
	Missing	958	
<b>Crowding</b>			
Number of children in house (6 months)	0	2595 (45)	2926 (44)
	1	2177 (38)	2454 (37)
	>2	1035 (18)	1180 (18)
	Missing	753	

Number of children in house (81 months)	0 1 >2 Missing	546 (10) 2888 (55) 1865 (35) 1261	673 (10) 3547 (54) 2340 (36)
Crowding index (8 months) <sup>5</sup>	≤0.5 0.5-0.75 0.75-1 >1 Missing	837 (16) 2192 (41) 1811 (34) 558 (10) 1162	1022 (16) 2479 (38) 2161 (33) 898 (13)
Crowding index (33 months) <sup>5</sup>	≤0.5 0.5-0.75 0.75-1 >1 Missing	1042 (20) 2042 (40) 1777 (34) 315 (6) 1384	1269 (20) 2474 (38) 2370 (36) 447 (7)
Crowding index (81 months) <sup>5</sup>	≤0.5 0.5-0.75 0.75-1 >1 Missing	712 (14) 1838 (36) 2123 (41) 501 (10) 1386	887 (14) 2206 (34) 2578 (39) 889 (13)
Presence of mould in the home (8 months)	None Not serious Fairly / Very serious Missing	4097 (75) 1236 (23) 143 (3) 1084	4912 (75) 1465 (22) 183 (3)
Presence of damp in the home (8 months)	None Not serious Fairly / Very serious Missing	2850 (52) 2356 (43) 289 (5) 1065	3196 (49) 2917 (45) 447 (6)
<b>Smoking</b>			

Child any exposure to smoke (up to 54 months) <sup>6</sup>	None Yes Missing	2654 (51) 2552 (49) 1354	3260 (50) 3300 (50)
Child exposure to smoke (6 months)	None Yes Missing	3845 (67) 1876 (33) 839	4331 (66) 2229 (34)
Child exposure to smoke (38 months)	None Yes Missing	3505 (63) 2033 (37) 1022	4209 (62) 2351 (38)
Child exposure to smoke (54 months)	None Yes Missing	3575 (65) 1912 (35) 1073	4306 (66) 2254 (34)
Maternal smoking during pregnancy (at any time) <sup>7</sup>	None Yes Missing	4730 (82) 1076 (19) 754	5310 (81) 1250 (19)
Maternal smoking during pregnancy (18 weeks)	None Yes Missing	4747 (83) 989 (17) 824	5374 (82) 1186 (18)
Maternal smoking during pregnancy (32 weeks)	None Yes Missing	4829 (87) 744 (13) 987	5631 (85) 929 (14)
<b>At birth</b>			
Breast feeding at any time point <sup>^</sup>	No Yes Missing	977 (16) 5019 (84) 564	1129 (17) 5431 (83)
Breast feeding duration across time points <sup>^</sup>	Never <3 months 3-5 months 6 months + Missing	1164 (20) 1327 (23) 977 (17) 2228 (39) 864	1419 (22) 1542 (24) 1123 (17) 2476 (38)

Birthweight of child	> 2500g	5832 (95)	6215 (95)
	≤ 2500g	311 (5)	345 (5)
	Missing	417	
Gestational age of child	< 37 weeks	339 (6)	359 (5)
	≥ 37 weeks	5872 (95)	6201 (95)
	Missing	349	
Child ever use a dummy/ pacifier	No	3160 (54)	3474 (53)
	Yes	2683 (46)	3086 (47)
	Missing	717	

\* Unless otherwise stated

\*\* All variables are derived from questionnaire data reported by the parent(s) when the child was 6, 18, 30, 42, 57, 69, 81, 91 and 103 months.

<sup>1</sup> A child was deemed to react to an allergen if the weal and/or flare was ≥2mm at the age 7 Focus clinic. Children were excluded if they didn't react to the positive allergen, or did react to the negative control. For this study children were categorised with an allergy if they were positive for house dust mite, mixed grasses or cat. These excluded children were imputed along with the missing data.

<sup>2</sup> Categorised as present if >1 cm in diameter in any of the following areas: around the eyes, the sides or front of the neck, in front of the elbows, behind the knees or in front of the ankles

<sup>3</sup> The child attended day care, as reported by the parent, at 15, 24 and 38 months.

<sup>4</sup> 'Regular' was defined as general habit and not a one off.

<sup>5</sup> A measure of the number of persons in the household divided by the number of rooms available (including the kitchen if large enough to eat in). The crowding index was derived from the number of persons in the household divided by the number of rooms available in the house (<0.5, 0.5-<0.75, 0.75-<1 or ≥1), where the higher the number the greater the crowding.

<sup>6</sup> Child had some exposure to smoking (passive smoking) at any point up to 54 months.

<sup>7</sup> Mother reported smoking during pregnancy (at any time).

<sup>^</sup> Owing to a difference in time points, numbers may differ.



Supplementary Table C. Fully adjusted model (original plus imputed data n= 6560).

		n (%)		Adjusted <i>p</i> value	Adjusted OR (95% CI)
<b>Model 1</b>		<b>No COM (n=5814)</b>	<b>Mild / significant COM (n=746)</b>	<b>No COM vs. mild / significant COM</b>	
Pus/mucus leaking from the child's ear	Never	4439 (76)	404 (54)	<b>&lt; 0.001</b>	<b>1.4 (1.3-1.6)</b>
	At least once	1375 (24)	342 (46)		
Child snores on at least two time points (prolonged snoring)*	No	5277 (91)	625 (84)	0.242	1.2 (0.9-1.5)
	Yes	537 (9)	121 (16)		
Child had grommets inserted (by age 9)	No	5634 (97)	563 (75)	<b>&lt; 0.001</b>	<b>4.9 (3.6-6.7)</b>
	Yes	180 (3)	183 (25)		
Child reported having tonsils removed (age 9)	No	5738 (99)	707 (95)	0.286	0.7 (0.4-1.3)
	Yes	76 (1)	39 (5)		
Child reported having adenoids removed (age 9)	No	5692 (98)	660 (88)	0.418	1.2 (0.8-1.9)
	Yes	122 (2)	86 (12)		
Tympanometry at age 7 (better functioning ear)	Type A <sup>R</sup>	4464 (77)	344 (46)	<b>&lt; 0.001</b>	<b>2.3 (1.9-2.9)</b>
	Type C1 / C2	1037 (18)	220 (29)		
	Type B	313 (5)	183 (25)		
Audiometry age 7 (better hearing ear)	< 20 dB	5694 (98)	696 (93)	0.996	1.0 (0.6-1.8)
	≥ 20 dB	120 (2)	50 (7)		
Birthweight	> 2500g	5524 (95)	691 (93)	0.463	1.2 (0.8-1.4)
	≤ 2500g	290 (5)	55 (7)		
Gestational age	≥ 37 weeks	5514 (95)	687 (92)	<b>0.010</b>	<b>1.5 (1.1-2.1)</b>
	< 37 weeks	300 (5)	59 (8)		

\*Because of coloration, 'prolonged snoring' was chosen to be entered in to the model over 'snoring' and 'early snoring' due to a higher univariate OR (odds ratio).

<sup>R</sup> Reference category

COM = chronic otitis media

CI = confidence interval